

Problem Set 6
CSCE 440/640

Due dates: Electronic submission of the pdf file of this homework is due on **11/2/2016 before 2:50pm** on ecampus.tamu.edu, a signed paper copy of the pdf file is due on **11/2/2016** at the beginning of class.

Name: (put your name here)

Resources. (All people, books, articles, web pages, etc. that have been consulted when producing your answers to this homework)

On my honor, as an Aggie, I have neither given nor received any unauthorized aid on any portion of the academic work included in this assignment. Furthermore, I have disclosed all resources (people, books, web sites, etc.) that have been used to prepare this homework.

Signature: _____

Problem 1. (20 points) Consider the mixed state

$$M = \left\{ \left(|0\rangle, \frac{1}{3} \right), \left(|1\rangle, \frac{2}{3} \right) \right\}.$$

- (a) Determine the density matrix ρ of the mixed state M .
- (b) Derive a different mixed state M' (which should not consist of computational basis states) that has the same density matrix ρ as M .

[This problem shows that density matrices are not in one-to-one correspondence with mixed states.]

Solution.

Problem 2. (20 points)

- (a) Do Exercise 3.5.1 (b) on page 55 of our textbook KLM.
- (b) Do Exercise 3.5.1 (c) on page 55 of our textbook KLM.

Solution.

Problem 3. (20 points) Find the Schmidt decomposition of the states

- (a) $\frac{1}{2} (|00\rangle + |01\rangle + |10\rangle + |11\rangle)$.
- (b) $\frac{1}{\sqrt{3}} (|00\rangle + |01\rangle + |10\rangle)$.

[Students of CSCE 440 only need to solve (a), and students of CSCE 640 should solve both (a) and (b).]

Solution.

Problem 4. (20 points) Exercise 3.5.4 (a) on page 57 in our textbook KLM.

Solution.

Problem 5. (20 points) Choi has shown that for all matrices $V_j \in \mathbf{C}^{n \times m}$, the map $T: M_n(\mathbf{C}) \rightarrow M_m(\mathbf{C})$ given by

$$T(\rho) = \sum_{j=1}^{\ell} V_j^* \rho V_j$$

is completely positive. Show that if the matrices V_j satisfy the condition

$$\sum_{j=1}^{\ell} V_j V_j^* = I,$$

where I denotes the identity matrix, then T is trace preserving, so $\text{tr } T(A) = \text{tr } A$. [Hint: the matrix trace satisfies $\text{tr}(ABC) = \text{tr}(CAB)$.]

Solution.

Checklist:

- Did you add your name?
- Did you disclose all resources that you have used?
(This includes all people, books, websites, etc. that you have consulted)
- Did you sign that you followed the Aggie honor code?
- Did you solve all problems?
- Did you submit the pdf file resulting from your latex source file on ecampus?
- Did you submit a hardcopy of the pdf file in class?