

Skip Lists

Andreas Klappenecker

Texas A&M University

© 2018 by Andreas Klappenecker. All rights reserved.

Goal

Consider a set

$$S = \{x_1 < x_2 < \dots < x_n\}$$

from a totally ordered universe. This set can dynamically change by adding or removing elements. Our goal is to search S for an element k .

Goal

Consider a set

$$S = \{x_1 < x_2 < \dots < x_n\}$$

from a totally ordered universe. This set can dynamically change by adding or removing elements. Our goal is to search S for an element k .

Bottom and Top Elements

We add a bottom element $-\infty$ and top element $+\infty$ to the universe such that

$$-\infty < x_1 < x_2 < \dots < x_n < +\infty.$$

These elements can simplify the implementation of the search.

Linked List Representation

Implementation

We can represent the set S by an ordered linked list. The problem is that we cannot index into this list, so the search is slow.

Search Trees

A search tree can speed up the search, but can be a bit awkward to maintain under insert and delete operations.

Idea Behind Skip Lists

We want to obtain the speed of a binary search tree but combine it with the ease of maintaining a sorted linked list.

Skip lists were invented by Bill Pugh in 1990.

They offer an expected search time of $O(\log n)$.

They generalize linked lists and are easy to implement.


A **descending filtration** is a sequence S_i of subsets of S such that

$$\emptyset = S_r \subseteq S_{r-1} \subseteq \cdots \subseteq S_1 = S.$$

In computer science, the S_i are called levels. The idea is that S_k for a large k is easy to search, since it has fewer elements than S_1 .

The idea is that we implement each S_i by a sorted linked list. Each element x in S_i is also linked to the element x in the finer level S_{i-1} .

Example



Search


When we search for k :

If $k = \text{key}$, done!


If $k < \text{next key}$, then k is not in this list, so go down a level

If $k \geq \text{next key}$, then go right

Example: Search for 61



Example: Search for 29



Construction

If the set $S_1 = S$ is fixed, then we could choose to include every other element into S_2 . Next, put every other element of S_2 into S_3 , and so forth.

Problem

We want to be able to insert and delete elements. These operations destroy the nice structure!

Construction

Let $S_1 = S$. For every element x in S_k , include x in S_{k+1} with probability $1/2$.

Expected Number of Elements

$$E[|S_1|] = n,$$

$$E[|S_2|] = n/2,$$

$$E[|S_3|] = n/4,$$

$$\vdots$$

We say that an element x_k has **height** ℓ if and only if

$$x_k \in S_\ell, \quad \text{but} \quad x_k \notin S_{\ell+1}.$$

Let X_k be the random variable that gives the height of the element x_k . We have

$$\Pr[X_k = \ell] = p(1 - p)^{\ell-1}.$$

So for $p = 1/2$, we have

$$\Pr[X_k = \ell] = (1/2)^\ell = 2^{-\ell}.$$

Interlude: Jensen's Inequality

Jensen's Inequality for Convex Functions

Proposition (Jensen's Inequality)

Let X be a random variable with $E[X] < \infty$. If $f: \mathbf{R} \rightarrow \mathbf{R}$ is a convex function, so \smile , then

$$f(E[X]) \leq E[f(X)].$$

Jensen's Inequality for Convex Functions

Proposition (Jensen's Inequality)

Let X be a random variable with $E[X] < \infty$. If $f: \mathbf{R} \rightarrow \mathbf{R}$ is a convex function, so \smile , then

$$f(E[X]) \leq E[f(X)].$$

Proof.

Since f is convex, we can find a linear function $g(x) = ax + b$ which lies entirely below the graph of f , but touches f at $E[X]$. In other words, we can choose real numbers a and b such that

$$f(E[X]) = g(E[X])$$

and $g(x) \leq f(x)$ for all $x \in \mathbf{R}$.

Proof. (Continued).

Since $f(x) \geq g(x)$ for all $x \in \mathbf{R}$, it follows that

$$\begin{aligned} E[f(X)] &\geq E[g(X)] \\ &= E[aX + b] = aE[X] + b \\ &= g(E[X]) = f(E[X]). \end{aligned}$$



Jensen's Inequality for Concave Functions

Proposition (Jensen's Inequality)

Let X be a random variable with $E[X] < \infty$. If $f: \mathbf{R} \rightarrow \mathbf{R}$ is a concave function, so \curvearrowright , then

$$E[f(X)] \leq f(E[X]).$$

Proof.

If f is concave, then $-f$ is convex. So

$$-f(E[X]) \leq E[-f(X)] = -E[f(X)]$$

by Jensen's inequality for convex functions. Thus,

$$f(E[X]) \geq E[f(X)]. \quad \square$$

Back to Skip Lists

Proposition

The expected maximum height of a skip list with n elements is given by

$$E \left[\max_{1 \leq k \leq n} X_k \right] \in O(\log n).$$

Proof.

Let α be a real number in the range $1 < \alpha < 2$. Then

$$\begin{aligned} \mathbb{E} \left[\max_{1 \leq k \leq n} X_k \right] &\leq \log_{\alpha} \mathbb{E} \left[\alpha^{\max_{1 \leq k \leq n} X_k} \right] \\ &= \log_{\alpha} \mathbb{E} \left[\max_{1 \leq k \leq n} \alpha^{X_k} \right]. \end{aligned}$$

Since $\alpha^{X_k} \geq 1$, we can estimate the right-hand side by the sum

$$\mathbb{E} \left[\max_{1 \leq k \leq n} X_k \right] \leq \log_{\alpha} \mathbb{E} \left[\sum_{k=1}^n \alpha^{X_k} \right].$$

Continued.

$$\begin{aligned} \mathbb{E} \left[\max_{1 \leq k \leq n} X_k \right] &\leq \log_{\alpha} \mathbb{E} \left[\sum_{k=1}^n \alpha^{X_k} \right] = \log_{\alpha} \left(\sum_{k=1}^n \sum_{k \geq 1} \alpha^k 2^{-k} \right) \\ &= \log_{\alpha} \left(\sum_{k=1}^n \frac{1}{1 - \alpha/2} \right) \\ &= \log_{\alpha} n + \log_{\alpha} \frac{1}{1 - \alpha/2} = O(\log n), \end{aligned}$$

which is what we wanted to show. □

Proposition

The number of levels of a skip list of a set with n elements satisfies $O(\log n)$ with high probability.

Proof.

Let X_k denote the random variable giving the number of levels of the k -th element of S . Then

$$\Pr[X_k > t] \leq (1 - p)^t.$$

So

$$\Pr[\max_k X_k > t] \leq n(1 - p)^t = \frac{n}{2^t}$$

for $p = 1/2$. Choosing $t = a \log n$ and $r = \max_k X_k$, we can conclude that

$$\Pr[r > a \log n] \leq \frac{1}{n^{a-1}}$$

for any $a > 1$. □